

CORRECTION

Open Access

Analysis of conditional gene deletion using probe based Real-Time PCR

Britta Weis^{1*}, Joachim Schmidt², Frank Lyko¹, Heinz G Linhart³

Abstract

Following publication of this article [1] the authors noticed that an incorrect probe reference was cited on page 3, 4, 5 and 6 ("UP #69, Roche Applied Science"). The correct probe that was used for the 1lox/2lox allele ratio analysis in the paper is as follows

Probe for 1lox/2lox allele quantification:

5'-**6-FAM-atAaCtTCgtatagCATaCattatac-BHQ-1** -3'

(uppercase letters = LNA bases)

Manufacturer: EUROGENTEC, Seraing, Belgium

All other information and reaction conditions in the paper are correct as stated.

Author details

¹Division of Epigenetics, German Cancer Research Center, Heidelberg, Germany. ²Division of Translational Oncology, German Cancer Research Center, Heidelberg, Germany. ³Department of Gastroenterology and Hepatology, University Hospital Freiburg, Hugstetter Str. 49, 79106 Freiburg, Germany.

Received: 13 December 2010 Accepted: 21 December 2010

Published: 21 December 2010

Reference

1. Weis B, Schmidt J, Lyko F, Linhart HG: **Analysis of conditional gene deletion using probe based Real-Time PCR.** *BMC Biotechnology* 2010, **10**:75.

doi:10.1186/1472-6750-10-90

Cite this article as: Weis et al.: Analysis of conditional gene deletion using probe based Real-Time PCR. *BMC Biotechnology* 2010 **10**:90.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed Central**

* Correspondence: b.weis@Dkfz-Heidelberg.de

¹Division of Epigenetics, German Cancer Research Center, Heidelberg, Germany

Full list of author information is available at the end of the article